



Research Article

Phytochemical study of *Aphanamixis polystachya* wall. root bark extract in various solvents

Lad Meenal^{1,*}, Sonar Shivani², Bhor Ila³
Professor & Head¹, P. G. Scholar², Associate Professor³

Department of Dravyaguna

PDEA's College of Ayurved and Research Centre, Nigdi, Pune, Maharashtra, India-411044

Article Received on: 03/01/2021; Accepted on: 17/02/2021

*Corresponding Author: Dr. Lad Meenal, E-mail: drmdlad@gmail.com

ABSTRACT:

In this present study photochemistry of dried root bark of *Aphanamixis polystachya* Wall has been done. The trial drug extract in various solvent such as water, PET ether, ethyl acetate, ethyl alcohol, methyl alcohol, chloroform in soxhlet apparatus were used for phytochemical investigation. As end result the different phyto-constituents obtained in extracts of various solvent are phenyl compounds, coumarins, fat and essential oils, terpenoids, monoterpenoids, diterpenoids, triterpenoids, steroids, alkaloids etc.

KEY WORDS: *Aphanamixis polystachya* Wall., Phytochemistry, phyto- constituents.

INTRODUCTION:

Different medical plants have been used for years in daily life to treat disease all over the world. Plants play a vital role in human and animal life as source of food and medicine. *Aphanamixis polystachya* shows many therapeutic properties as cardiogenic and choleretic activity [1], antioxidant [2]. *Amoora rohituka* is described in Ayurveda, an Indian traditional system of medicine for management of disorders of blood, diseases of eye, helminthiasis disease, ulcer, liver disorders and splenomegaly. Researchers studied that many medicinal plants possess significant hepatoprotective activity [3]. The fruit, bark and leaves extracts and of *Aphanamixis polystachya* possess significant cytotoxic and antioxidant activity, thrombolytic, antimicrobial activity [4,5]. Limonoids from *Aphanamixis polystachya* shows Antifeedant Activity [6]

Aphanamixis polystachya pacifies vitiated vata, and pitta, splenomegaly, liver disorders, tumor, ulcer, dyspepsia, intestinal worms, skin diseases, diabetes, eye diseases, jaundice, hemorrhoids, burning sensation, rheumatoid

arthritis and leucorrhoea. [7]. The extract of stem bark, leaves and leaf derived callus of *Amoora rohituka* in various solvent were examined against 10 species of human pathogenic bacteria. [8] Researcher evaluate *in vitro* antibacterial activity, cytotoxic activity and *in vivo* anti-diarrheal activity of ethanolic extract of *Aphanamixis polystachya* (wall.) Parker leaf and concluded that Leaf extract of *Aphanamixis polystachya* (wall.) Parker to be a good source of antimicrobial and cytotoxic property. [9]. The study is done on characterization and fatty acid composition of *Amoora rohituka* Seed and Leaf Oils and concluded that both seed and leaf oils of *Amoora rohituka* are rich in essential fatty acids and consequently the oils especially the seed oil may prove its worth as edible oil if considered from nutritional point of view [10]. The extracts of *Amoora rohituka* leaf were tested against the red flour beetle and investigate the biological activity such as dose-mortality, insect repellency, cytotoxicity, larvicidal and antimicrobial assays [11]. Researcher investigate the crude n-hexane, ethyl acetate and

methanol extracts of *Aphanamixis polystachya* leaves for their antimicrobial, antioxidant, cytotoxic and thrombolytic activities. [12]

After review of literature the detail study of phytochemical studies of *Aphanamixis polystachya* Wall. root bark under identical set of experimental condition is still lacking. It was thought of interest to study the phyto-constituents present in *Aphanamixis polystachya* Wall. Root bark under suitable condition.

Aim:

To study phytochemicals or phyto-constituents present in *Aphanamixis polystachya* wall. root bark extract in various solvents.

MATERIALS:

Aphanamixis polystachya root bark was collected from the garden of College of Ayurved and Research Centre, Nigdi, Pune (India) in the month of April 2019 at morning time. Chloroform, Ethanol, Ethyl acetate, n-Hexane, Methanol, Petroleum ether are used as solvents for extraction. Mayer reagent, Wagner's reagent, Hager's reagent, Dragendroff's reagent, Lead acetate, Ferric chloride, Potassium dichromate, Sodium

hydroxide, Sodium chloride, gelatin solution, Ammonia solution, Zinc chloride, Hydrochloric acid, Glacial acetic acid, Molisch's reagent, copper acetate, Glacial acetic acid, Cobalt chloride, Benedict's reagent, Fehling solution A and B, Ninhydrin, Million's reagent, Sodium nitroprusside, Sulphuric acid, Nitric acid, Iodine, Safranin, picric acid, potassium hydroxide, Ammonium hydroxide, benzene, pyridine, gallic acid, All the chemicals were used of AR grade (Sd fine chemicals Pvt Ltd., Mumbai).

Preparation of extracts:

The air-dried root bark of *aphanamixis polystachya* (wall) *parker* was shade dried and powdered at room temperature. 10g of powder was subjected to successive hot continuous extraction (soxhlet) in various solvents.

METHODS:

Collection of root-bark of *Aphanamixis polystachya* is collected from matured and well developed tree. Bark of perennial root is collected in early spring (March). Collected material is washed thoroughly and dried.[13]

The phytochemical investigation were done by using known methods [14-20]

Observations and Results:

Table No. 1: Phytochemical screening of *Aphanamixis polystachya* root bark in various solvent.

Phytochemicals	Water	PET ether	Ethyl acetate	Methyl alcohol	chloroform	Ethyl alcohol
Alkaloids	+	+	+	+	+	+
Phenolic compound	-	-	-	-	-	+
Flavonoides	+	-	+	+	-	+
Carbohydrates	-	-	+	+	-	+
Reducing sugar	+	-	+	+	-	+
Amino acid	-	-	-	-	-	-
Phytosterol	+	-	+	+	-	+
Protein	-	-	-	+	-	-
Gum and mucilage	-	-	-	-	-	-
Anthocyanin	-	-	-	+	-	+
Glycoside	-	-	-	+	-	-
Saponin	+	+	+	-	+	-
Coumarin	-	-	+	+	-	+
Emodine	-	-	-	-	-	-
Adonine	-	-	-	-	-	-
Diterpines	+	-	+	-	-	+
Cardiac glycoside	-	-	+	+	-	+
Tannin	-	-	-	+	+	-

The result of phytochemical screening of *Aphanamixis polystachya* in various solvents shows in table. Extract of water shows presence of alkaloids, flavonoids, reducing sugar, phytosterols, diterpines and saponin. Extract of PET ether shows presence of alkaloids and saponins. Extract of ethyl acetate shows presence of alkaloids, flavonoides, carbohydrates, reducing sugar, phytosterol, saponin, coumarin, diterpines, and cardiac glycoside. Extract of methyl alcohol shows presence of alkaloids, flavonoides, carbohydrates, reducing sugar, phytosterol, protein, anthocyanin, glucoside, saponin, cardiac glycoside and tannin. Extract of chloroform shows presence of alkaloids, glycoside and tannins. Extract of ethyl alcohol shows presence of alkaloides, phenolic compound, flavonoides, carbohydrates, reducing sugar, phytosterol, anthocyanin, coumarin, diterpines and cardiac glycoside.

DISCUSSION:

Phytochemical analysis is very important to rule out adulteration in the herbal formulations. Eg. Root bark of Rohitaka is useful in treatment but due to non-availability the use of stem bark is regular in routine practice. There are several species available in different floras and in market trade. Root bark of *Aphanamixis polystachya* contains phyto-constituents such as alkaloids, flavonoids, reducing sugar, phytosterols, carbohydrates, diterpines and saponin. Alkaloids shows analgesic activity, which is homologous to Vataghna activity which is useful in conditions like arthritis. Terpenoids have medicinal properties as analgesic and antiseptic which is same as Vataghna and Vranashoshana activity respectively. These activities are useful for treating septic wound. Cardiac glycosides shows cardio-tonic activity which is same as Hridya activity which can be used in heart disease. This study can be further continued for finger printing of phyto-constituents present in *Aphanamixis polystachya*.

Future Scope-

Comparative phytochemical study if root-bark and stem-bark of *Aphanamixis polystachya* can be done in future.

CONCLUSION:

This study can be concluded as the phytochemical analysis in various solvents of root bark of *Aphanamixis polystachya* shows the presence of Alkaloids, Flavonoids, Phytosterols, Caumarine, Diterpines, Cardiac Glycosides.

REFERENCES:

- [1] Raihan Khan Tanveer, Karmarkar Palash, Das Abhijit, Banik Rana, Sattar Md. Mafruhi, Evaluation of cytotoxic and anthelmintic activities of bark extract of *Aphanamixis polystachya*(wall) *Int. Res. J. Pharm.*, 4(40):126-128, 2013.
- [2] Kavitharaj V, Avinash K O, Kumar J R, Murugesan Ashok, Gnanasekaran, Karthikey and Chandrashekrappa GK, In vitro antioxidant activities of stem bark, leaves and callus extracts from *Amoora rohituka*, *International Journal of Phytomedicine* 9:584-588, 2017.
- [3] Vuyyuri Bhaargavi, G.S.L. Jyotsna and Reshma Tripurana, A review on hepatoprotective activity, *IJPSR*, Vol. 5(3): 690-70., 2014.
- [4] Apurba Sarker Apu, Fariha Akhter Chowdhury, Farjana Khatun, A.T.M. Jamaluddin, Atiqul Haque Pathan and Arindom Pal, Phytochemical Screening and *In vitro* Evaluation of Pharmacological Activities of *Aphanamixis polystachya* (Wall) Parker Fruit Extracts, *Tropical Journal of Pharmaceutical Research* February, 12(1): 111-116, 2013.
- [5] Nahid sultana, M. Oliur Rahman, Faizatahia, Md. Abul Hassan and Md.R Rashid, antioxidant, cytotoxicity and antimicrobial activities of *Aphanamixis polystachya* (wall.) R. N. parkar, Bangladesh, *J. Bot.*, 46 (4): 1381-1387, 2017.
- [6] Jie-Yun Cai, Duo-Zhi Chen, Shi-Hong Luo, Ning-Chuan Kong, Yu Zhang, Ying-Tong Di, Qiang Zhang, Juan Hua, Shu-Xi Jing, Shun-Lin Li, Sheng-Hong Li, Xiao-Jiang Hao and Hong-Ping He, Limonoids from *Aphanamixis polystachya* and their Antifeedant Activity, *J. Nat. Prod.*, 77: 472-482, 2014.
- [7] Malviya S, Jain S, Patel R, Malviya V. Pharmacological review on *Amoora rohituka*, *World J Pharm Pharm Sci*, 1(1): 7-16, 2012.

- [8] Ashwini H.M. and Ravishankar Rai V , In vitro evaluation of the Efficacy of stem bark , leaves and callus extract of *Amoora rohituka* on important human pathogenic bacteria. , *Indian Journal of Applied Microbiology*,10(1): 88-92, 2009.
- [9] H. M. Shadid Hossain Snigdha, Rezwana Ali, Dipankar Kumar Das and Md. Abdul Wadud, Biological evaluation of ethanolic extract of *Aphanamixis polystachya* (Wall.) Parker leaf, *Int. J. Adv. Multidiscip. Res.*, 3 (9): 13-21, 2016.
- [10] Moumita Gupta, Pijush Kundu and Subrata Laskar, Characterization and Fatty Acid Composition of *Amoora rohituka* Seed and Leaf Oils, *Biosciences, Biotechnology Research Asia* ,Vol. 8(1), 213-218, 2011.
- [11] Md. Rubel Rana, Nowrin Islam Amin, Abdullah An Naser and Nurul Islam, Biological activities of *Amoora rohituka* Roxb. Leaf extracts through dose-mortality, insect repellency, cytotoxicity, larvicidal and antimicrobial assays, *Journal of Pharmacognosy and Phytochemistry* , 6(3): 606-611, 2017.
- [12] Apurba Sarker Apu, Atiqul Haque Pathan, Abu Taiab Md. Jamaluddin, Ferdous Ara, Shakhawat Hossan Bhuyan and Md. Rakibul Islam, Phytochemical Analysis and Bioactivities of *Aphanamixis polystachya* (Wall.) R. Parker Leaves from Bangladesh, *Journal of Biological Sciences*,13 (5): 393-399 , 2013.
- [13] Guidelines On Good Field Collection Practices For Indian Medicinal Plants, National Medicinal Plant Board, Department of AYUSH, 17, 2009.
- [14] K. Santhi and R. Sengottuvel, Qualitative and Quantitative Phytochemical analysis of *Moringa concanensis* Nimmo, *Int.J.Curr.Microbiol.App.Sci*, 5(1): 633-640 , 2016.
- [15] Teresa May B. Bandiola, Extraction and Qualitative Phytochemical Screening of Medicinal Plants: A Brief Summary, *Int J Pharm* , 8(1): 137-143, 2018.
- [16] Prashant Tiwari, Bimlesh Kumar, Mandeep Kaur, Gurpreet Kaur, Harleen Kaur, Phytochemical screening and Extraction: A Review, *Int.Vol.1(1):98-106*,2011.
- [17] Sarla Saklani, Abhay P. Mishra, Bhawana sati, Hemlata sati, Pharmacognostic, phytochemical and microbial screening of *Aphanamixis polystachya* , an endangered medicinal tree , *Int J Pharm Pharm Sci*, Vol 4, Suppl 3: 235-240 ,2012.
- [18] Ejoba Raphael, Phytochemical constituents of some leaves extract of *Aloe vera* and *Azadirachta indica* plant species, *Global Advanced Research Journal of Environmental Science and Toxicology*, Vol. 1(2) pp. 014-017, 2012.
- [19] Shweta S. Saboo, Rani W. Chavan, G.G.Tapadiya1, S.S.Khadabadi, Detail phytochemical study of *Aphanamixis Polystachya* (wall) parker, *world j. of pharmacy and pharmaceutiocal sci.*, Volume 3, Issue 3:1583-1594.,2014.
- [20] S Sridvidya , G Sridevi , A G Manimegalai, Phytochemical Screening and In Vitro Antioxidant Activity of Ethanolic Extract of *Cassia occidentalis*, *International Journal of Pharmaceutical and Clinical Research* , 9 (3): 252-256, 2017.
- [21] Gibbs R.D., (1974) Chemotaxonomy of Flowering Plants. Vol.1, McGill Queen's University Press, Montreal and London., Narasinga Rao V and DSVGK Kaladhar, Phytochemical and biochemical studies of medicinal plant *Globba Bulbifera*, *Inter. J. of Phytotherapy*, Vol 4 (1): 50-53, 2014.

Cite this article as:

Lad Meenal, Sonar Shivani, Bhor Ila, Phytochemical study of *Aphanamixis polystachya* wall. root bark extract in various solvents, *PDEAS Int. J. R. Ayu. & A. Sc.*, 3(1); Feb 2021, p. 01-04.